

ISBN : 978-979-96595-4-5



Proceeding

The 2nd International Conference of
the Indonesian Chemical Society 2013

IC  CS 2013

Research in Chemistry for Better Quality of Environmental

Universitas Islam Indonesia, Yogyakarta, Indonesia
October, 22 - 23rd 2013

Abdul Kahar Muzakkir, Conference Hall
Universitas Islam Indonesia (UII), Yogyakarta.
Kampus Terpadu, Jl. Kaliurang KM 14,5 Sleman, Yogyakarta.

Preface

The international conference is an annual conference of the Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI). In the year 2013, the mandate of the organizing committee was given to the HKI Yogyakarta branch and also supported by Department of Chemistry of Universitas Negeri Yogyakarta (UNY), Department of Chemistry of Universitas Gadjah Mada (UGM), Department of Chemistry of Universitas Islam Negeri Sunan Kalijaga (UIN Suka), National Nuclear Energy Agency (BATAN Yogyakarta), and Volcano Investigation and Technological Development Center (BPPTK Yogyakarta). For the year 2013, ICICS 2013 is hosted by Department of Chemistry, Faculty of Mathematics and Natural Sciences, Islamic University of Indonesia, Yogyakarta from October 22 – 23, 2013. This conference was also prepared to celebrate 70th anniversary of Universitas Islam Indonesia.

The Scientific Programme of ICICS2013 comprises the following:

- | | | |
|---|----|--------|
| 1. Invited Speaker | 11 | papers |
| 2. A total 256 paper for parallels sessions | | |
| a. Organic Chemistry | 32 | papers |
| b. Inorganic Chemistry | 43 | papers |
| c. Physical Chemistry | 37 | papers |
| d. Analytical Chemistry | 68 | papers |
| e. Education Chemistry | 23 | papers |
| f. Biochemistry | 43 | papers |

The breakdown of the presentation is as follows:

Session	Oral	Poster	Total
Invited Speaker	11	0	11
Organic Chemistry	25	7	32
Inorganic Chemistry	38	5	43
Physical Chemistry	31	6	37
Analytical Chemistry	61	7	68
Education Chemistry	22	1	23
Biochemistry	34	8	43
Total	222	34	256

Yogyakarta, 25th November 2013



Editors

Welcoming Address by The Organizing Committee



Assalamu'alaikum Wr. Wb.

Honorable Rector of Universitas Islam Indonesia
The distinguished invited speakers, and
All participants of the ICICS 2013

Welcome you at the 2nd International Conference of the Indonesia Chemical Society 2013 (ICICS 2013) this morning here at the Auditorium Kahar Muzakkir Universitas Islam Indonesia, Yogyakarta. The international conference is an annual conference of the Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI). In the year 2013, the mandate of the organizing committee was given to the HKI Yogyakarta branch and also supported by Department of Chemistry of Universitas Negeri Yogyakarta (UNY), Department of Chemistry of Universitas Gadjah Mada (UGM), Department of Chemistry of Universitas Islam Negeri Sunan Kalijaga (UIN Suka), National Nuclear Energy Agency (BATAN Yogyakarta), and Balai Penyelidikan dan Pengembangan Kegunungapian (BPPTK Yogyakarta). For the year 2013, the honor of hosting ICICS 2013 has been given to the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia, Yogyakarta. This conference was also prepared to celebrate 70th anniversary of Universitas Islam Indonesia.

The conference comprises both oral and poster presentation in English and Indonesian with optional post conference publication of full papers in English in the *Procedia Chemistry* (Elsevier, ISSN: 1876-6196) and *Proceeding Conference for Indonesian language*. There are 211 papers presented orally and 34 papers presented by poster covering wide-variety subjects of chemistry. We invited 6 Indonesian invited speakers, 2 Japan invited speakers, 1 Australian invited speakers, 1 Saudi Arabia invited speakers, and 1 Malaysian Invited speakers.

We hope you will enjoy a pleasant and valuable seminar at Universitas Islam Indonesia

Wassalamu'alaikum Wr. Wb.

Riyanto, Ph.D.



Opening Speech from the Rector of Universitas Islam Indonesia



Assalamu'alaikum Wr. Wb
The distinguished invited speakers, and
All participants of the ICICS 2013

Firstly, I would like to express my great appreciation to the Department of Chemistry UII as one of the organizers of the program The 2nd International Conference of the Indonesian Chemical Society 2013 (ICICS 2013) with the theme "Research in Chemistry for Better Quality of Environmental". I am proud that this interesting event is being organized and held in Yogyakarta.

As the biggest and the oldest private university in Yogyakarta, University Islam Indonesia is committed to the excellence in research and teaching. Recently, we are preparing UII as one of the world class universities.

Knowing that committee has selected outstanding speakers from various prestigious institutions. I believe that all of the participants will enjoy the discussion of issue covered by the topic of this seminar. Scientist have shown that the environment's condition is increasingly critical, and human industrial activities are largely to blame. In fact that environmental damage is a crisis we caused together, therefore, a responsibility we all share together. We are deeply concerned with the issues and opportunities in the internationalization of sciences for better life, sciences have to make better quality of environmental.

Finally, I would once again like to thank the organizer for organizing this event, and to thank all the participants attending this ICICS 2013 event as well as delivering their scientific presentations. I do really hope that you can enjoy this seminar and have excellent stay in Yogyakarta.

Wassalamu'alaikum Wr. Wb

Prof. Dr. Edy Suandi Hamid, M.Ec.
Rector of Universitas Islam Indonesia

Remarks by the Chairman of the Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI)



Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI) is an independent, nonprofit organization founded in February 1962 to facilitate communication among Indonesian chemists and other professionals from chemistry related fields, and to promote the advancement of science, education, and application of chemistry to support the better life of mankind. HKI organize activities to enhance communication and collaboration among chemists in various institutions in Indonesia, to disseminate new knowledge and research results in chemistry and related fields, to improve the knowledge and

skills of chemists working in schools, universities, industries, research institutes, and other sectors, to nurture a scientific temper on school children to ensure strong capabilities of future chemists that are needed for humankind, and other activities that support its missions. HKI holds various academic conferences, publishes several journals, supports the development of scientific information systems in Indonesian; organize training for chemists in various sectors, etc.

The 2013 International Conference of the Indonesian Chemical Society will be the 2nd event in the ICICS conference series, started in 2012, that brings together individuals involved in chemistry-related fields (chemistry, pharmacy, environmental science, chemical engineering, molecular biology, material science, education chemistry, etc.) or institution in chemistry-related sectors. The First International Conference of the Indonesian Chemical Society 2012 is organized by East Java Branch of HKI in collaboration with chemistry departments at several universities in East Java: ITS, UB, UIN Maliki, UM, UMC, Unair, Unej, and Unesa.

ICICS 2013 will be organized by the Indonesian Chemical Society Yogyakarta branch. The international conference was supported by the Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI), Department of Chemistry of Universitas Negeri Yogyakarta (UNY), Department of Chemistry of Universitas Gadjah Mada (UGM) and Department of Chemistry of Universitas Islam Negeri Sunan Kalijaga (UIN Sunan Kalijaga). For the year 2013, the honor of hosting ICICS-2013 has been given to the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Islam Indonesia (UII), Yogyakarta, Indonesia.

Congratulations to the ICICS 2013 committee for this conference.

Dr. Muhamad Abdulkadir Martoprawiro
Chairman of the Indonesian Chemical Society

Committees

Steering Committee

1. Head of the Indonesian Chemical Society
2. Head of the Indonesian Chemical Society Yogyakarta Branch
3. Rector of Islamic University of Indonesia
4. Rector of Gadjah Mada University
5. Rector of Yogyakarta State University
6. Rector of State Islamic University of Sunan Kalijaga
7. Head of Geological Agency of Indonesia
8. Head of National Nuclear Energy Agency of Indonesia

Reviewers:

1. Prof. Dr. Hardjono Sastrohamidjojo (UII Yogyakarta)
2. Prof. Dr. Nurfina Aznam (UNY Yogyakarta)
3. Prof. Dr. Karna Wijaya (UGM Yogyakarta)
4. Prof. Harno Dwi Pranowo (UGM Yogyakarta)
5. Prof. Drs. Sahat Simbolon, M.Sc. (BATAN Yogyakarta)
6. Dr. Muhamad A. Martoprawiro (HKI)
7. Riyanto, Ph.D. (UII Yogyakarta)
8. Dr. Is Fatimah (UII Yogyakarta)

Editors:

1. Dr. Noor Fitri (UII Yogyakarta)
2. Drs. Allwar, M.Sc., Ph.D. (UII Yogyakarta)
3. Rudy Syah Putra, Ph.D. (UII Yogyakarta)
4. Dwiwarso Rubiyanto, M.Si. (UII Yogyakarta)
5. Tatang Shabur Julianto, M.Si. (UII Yogyakarta)

Proceeding

The 2nd International Conference of the Indonesian Chemical Society 2013
October, 22-23th 2013

Organizing Committee :

- Chairperson : Riyanto, Ph.D.
- Secretary : 1. Rudy Syahputra, Ph.D.
2. Cecep Sa'bana Rahmatillah, S.Si.
3. Siswanta, S. Pd.
- Treasure : 1. Dr. Noor Fitri, M.Si.
2. Masirah, A.Md
3. Yuni Ari Rahmawati, S.Ag.
- Programmer & protocol : 1. Sunaryo MD
2. Lindung Prasetya Kurniawan, A.Md.
3. Paidi
4. Asih
5. Vira Megasari Haqni, ST
- Proceedings : 1. Thorikul Huda, S.Si. M.Sc.
2. Krisna Merdekawati, S.Pd., M.Pd.
3. Puji Kurniawati, S.Pd.Si., M.Sc.
4. Bayu Wiyantoko, S.Si., M.Sc.
5. Tri Esti Purbaningtias, S.Si., M.Si.
6. Yuli Rohyami, S.Si., M.Sc.
7. Dedy Sugiarto, S.Si.
- Transportations and Accommodations : 1. Jamalul Lail, S.Si.
2. Ponimin, SE
3. Agus Sri Untoro
4. Sukadi
5. Parwanto
- Appurtenance : 1. Drs. Sunarwi
2. M. Achnaf, A.Md.
3. Sigit Mujiarto
4. Kuntoro Haryanto, A.Md.
5. Dwi Mahmudi, BA
- Logistic : 1. Reni Banowati Istiningrum, S.Si.
2. Indriyani, A.Md.
3. Painem
4. Syaida
5. Sukirman
6. Aprilia Risky Wijaya

Proceeding

The 2nd International Conference of the Indonesian Chemical Society 2013
October, 22-23th 2013

Publication and Documentation : 1. Anang Susilo, A.Md.
2. Sihono
3. Umar Hasyim
4. Christanto Yuwono
5. Suratmin

Supporting Team : Himpunan Mahasiswa Kimia (HMK UII)



CONTENT

Content		Page
Cover		i
Preface		ii
Welcoming address by The Organizing Committee		iii
Opening Speech from the Rector of Universitas Islam Indonesia		iv
Remarks by the Chairman of the Indonesian Chemical Society (Himpunan Kimia Indonesia, HKI)		v
Committe		vi
Reviewers and Editors		vi
Content		ix
Invited Speaker		
Shaobin Wang, Stacey Indrawirawan, Yunjin Yao, Hongqi Sun	Graphene Supported Oxide Systems for Catalytic Oxidation of Organic Compounds in Aqueous Solution for Water Treatment	xii
Tatsufumi Okino	Chemistry and biology of brominated compounds from marine algae <i>Laurencia</i> spp.	xv
Heriyanto, Leenawaty Limantara	Chlorophyll and Carotenoid Prospects on Food, Health and Energy	xviii
Katsumi Kaneko	Molecular Functions of 1 nm-Scale Pore Spaces and their Application Potential to Sustainable Technologies	xxviii
Fethi Kooli	Al ₁₃ Intercalated and Pillared Montmorillonites from Unusual Antiperspirant Aqueous Solutions: Precursors for Porous Clay Heterostructures and Heptane Hydro-Isomerization Catalytic Activities	xxxi
Allwar, Ahmad Md. Noor, Mohd Asri bin Mohd Nawi	Characterizing Microporous Structures using Nitrogen Adsorption- Desorption Isotherm for Activated Carbon Prepared with Different Zinc Chloride Concentrations	xxxv
Papers of Inorganic Chemistry		
Ahmad Budi Junaidi, Helda Rahmawati dan Utami irawati	Study of Carboxymethyl Chitosan Synthesis : Effect of NaOH Concentration and Rtio Chitosan/Monochloro Acetic Acid Toword On The Substitution Degree and Solubility In Water	1-5
Ahmad Suseno, Priyono, Karna Wijaya,Wega Trisunaryanti	Study of Structure and Morphology of Surfactant-Modified Al- pillared Natural Bentonite	6-14
Aman Sentosa Panggabean, Subur P. Pasaribu, Dadan Hamdani, Nadira ¹	Synthesis of A Chelating Resin Chitosan-1,5-Diphenyl Carbazide and Characterization of Retention toward Cr(VI) Ions	15-25
Anti K. Prodjosantoso	Preparation and Characterisation of Chloride-Free Palladium Catalysts	26-32

Proceeding

The 2nd International Conference of the Indonesian Chemical Society 2013
October, 22-23th 2013

Dwi Rasy Mujiyanti, Totok Wianto, M. Fahmi Arif	Synthesis and Characterization of Nanosilica from Rice Husk Ash by Sol-Gel Process	33-38
Endang Susilowati, Triyono, Sri Juari Santosa, Indriana Kartini	Synthesis of Silver-Chitosan Nanocomposites by Glucose As Reducing Agent and Their Antibacterial Activity	39-47
Febriyana Rizky Hapsari, Ersalina Nidianti, Warsito, Edi Priyo Utomo	Analysis In Silico on Structure-Odor Relationship (SOR) Of Organoleptic-Tested Compounds	48-55
Indah Uswatun Hasanah, Ria Armunanto, Bambang Setiadji	The Oretical Study Properties of Semiconductor Metalloporphyrin Complexes Calculated Density Functional Theory Method (DFT)	56-66
Jannatin 'Ardhuha	Characterization of The FePd/NiTi Shape Memory Alloy Film for Sensor Applications	67-76
Lia Destiarti, Nelly Wahyuni, Ahmad Yani	Synthesis of Zeolite A from Capkala Kaolin by Varying Mass of Alumunium Oxide: XRD spectrum and CEC number of products	77-82
Maula Eka Sriyana, Aang Hanafiah W.S.	Physicochemical Characteristics of ^{99m} Tc-dtpa-ketoconazole as A Radiopharmaceutical for Deep Seated Fungal Detection	83-89
Muhamad Basit Febrian, Aminudin Sulaeman, Muhayatun Santoso	Preliminary Study of Comparison of EDXRF and ICP-AES Techniques for the Measurement of Elements in Fine Particulate Matter (PM _{2.5}) : Accuracy and Precision of XRF Technique	90-94
Muhdarina, Nurhayati, Flora Sijabat	Characterization of Phosphated Palas Clay	95-100
Nugrahaning Wuri Hakiki, Maria Christina P., Isti Daruwati	Penandaan M41S-NH ₂ dengan Radionuklida Teknesium-99m : Perbandingan Metode Langsung dengan Metode Tidak Langsung dalam Aplikasi Radiosinovektomi	101-111
Nurul Hidayati Fithriyah, Erdawati	Preservation of Paper Samples Coated With Chitosan Nanoparticle	112-120
Putu Sukmabuana, Poppy Intan Tjahaja, and Anton Winarko	A Surveilence on Tritium Radionuclide in Surface Soil of The TRIGA 2000 Reactor Site, Bandung	121-128
Ria Armunanto[*], Karna Wijaya, Radite Yogaswara	Study of CO Adsorption on Ni _n ^q (n=3-5; q=0, 1, -1) Clusters using DFT Method	129-139
Restu Kartiko Widi, Arief Budhyantoro, Emma Savitri	Reaction Study of Phenol Hydroxylation on Al/Fe Pillared and HDTMA Intercalated Bentonite Catalyst	140-146
Singgih Hartanto, Achmadin Luthfi, Sri Handayani	Characterization of Membrane PVA/Silica and PVA/Zeolite for Purification Bioethanol by The Vapor Permeation Process.	147-152
Triastuti Sulistyaningsih, Sri Juari Santosa, Dwi Siswanta, Bambang Rusdiarso	Hydrothermal Efek on Magnetite-Mg/Al-NO ₃ -HT Composite Synthesis	153-160
Tutik Setianingsih, Indriana Kartini, Yateman Arryanto	Synthesis of Mesoporous Carbon from Fructose by using Activator of Zinc Borosilicate at Low Temperature	161-172
Linda Dwitasari, Tutik Dwi Wahyuningsih, Indriana Kartini	HOMO and LUMO Determination of Chlorophyllin and Xanthophyll Dyes using Cyclic Voltammetry	173-178
Erni Astuti, Yateman Arryanto, Indriana Kartini	Facile Hydrothermal Synthesis of Various Nanostructured Titania	179-184

Proceeding

The 2nd International Conference of the Indonesian Chemical Society 2013
October, 22-23th 2013

Arief Rahmatulloh, Lukman Atmaja, Nurul Widiastuti	Korelasi Konsentrasi <i>Silane</i> dan Suhu Operasi Terhadap Konduktivitas Membran Komposit Kitosan – <i>Fly Ash</i> untuk Aplikasi <i>Proton Exchange Membrane Fuel Cell</i>	185-195
Paulina Taba, Marthinus Pongsendana, Eldayanti Ruru	Thiol-Functionalized Mesoporous Silica, MCM-48 as Adsorbent Ag(I) and Cd(II) Ions	196-201
<u>F. Widhi Mahatmanti</u>, Nuryono, Narsito	Synthesis of Chitosan-Silica Film using Sodium Silicate Solution from Rice Hull Ash	202-206
Maria Dewi Astuti, Dwi Rasy Mujiyanti, Dahlena Ariyani, Mustika Rahmadini	Perbandingan Sifat Karakteristik Silika Gel Sintesis dari Abu Sekam Padi Daerah Gambut dan Komersial	207-212
Husna Amalya Melati	Corrosion Protection Efficiency of Hybrid Polymers Coatings based TMSPMA Monomers on Carbon Steel in Saline Environment Evaluated by Electrochemical Measurements	213-219

Reaction Study of Phenol Hydroxylation on Al/Fe Pillared and HDTMA Intercalated Bentonite Catalyst

Restu Kartiko Widi*, Arief Budhyantoro, Emma Savitri

Department of Chemical Engineering, University of Surabaya, TG 6th floor, Raya Kalirungkut, 60293,
Surabaya, Indonesia
*restu@ubaya.ac.id

Abstract

The phenol hydroxylation reaction over Al/Fe Pillared and HDTMA intercalated bentonite has been studied using batch system. The effects of the preparation methods on the catalytic performance of modified bentonite were investigated. Natural bentonite was modified by direct and indirect pillarisation. In direct method, Al and Fe ions were used as pillaring agent to produce a bigger clay pore size. In indirect method, the first step was intercalation process to insert organic surfactant molecules (HDTMA) inside the clay molecules, and then the next step was pillarisation process using Fe cationic. Pillarisation of natural bentonite and HDTMA-bentonite with Fe metal has been done in various mol Fe solutions. The ratio of bentonite and intercalating agent or pillaring agent was 1 gr/50 ml. The mixture was agitated, and then the solid phase was washed with distilled water until free of chlorine ion. Then it was dried and calcined at 450 °C for 6 hours in N₂ and O₂ atmosphere. The modified bentonites were characterized by FT-IR spectrometer.

The catalytic reaction was conducted by entering both catalyst and phenol into batch reactor and then the mixture was heated. The reaction was conducted at 70°C for 4 hours. Along reaction (after temperature 70°C reached), solution of H₂O₂ was added stepwise into reactor during reaction. Molar ratio of phenol to H₂O₂ is 1 : 1. The weight ratio of catalyst to phenol was 1:10, and concentration of phenol is 1 M. The observed variables were catalyst activity and selectivity. The catalyst activity was observed as phenol conversion, and the selectivity was observed as selectivity of wanted product (hydroquinone).

The presence of Fe-pillared bentonite catalyst in the phenol hydroxylation reaction will give effect on increasing catalyst activity. It happens because number of active site and suitable pore size increase so that the amount of phenol, which can be converted into product, will also increase. It gives impact on selectivity of hydroquinone, cathecol, and benzoquinone will also increase. Finally, as conclusion the optimum performance of catalyst is PILB HDTMA 0.1 % Fe 1 mmol in the phenol hydroxylation reaction.

Keyword: bentonite, pillarisation, intercalation, phenol hydroxylation, hydroquinone

Introduction

Clay is one of the most abundant material found in nature, especially in Indonesia. Bentonite, one of material kind of clay, which is used in this research, is natural bentonite from Tulakan, Pacitan in east Java. Bentonite is generated frequently from the alteration of volcanic ash, consisting predominantly of smectite minerals, usually montmorillonite (Na,Ca)_{0.33}(Al,Mg)₂Si₄O₁₀(OH)₂·nH₂O. Up to now, bentonite can be utilized as filler, ion exchanger, catalyst, adsorbent etc.

Bentonite has characteristics such as moisture content as high as 30-40 %, and easy to be swelling. Because of the properties, the potential of bentonite application has recently

attracted a lot of attention in both academia and industry, which concern the understanding of maximize the existing potential. The characteristics cause unstable structure and make low selectivity for both adsorbent and catalyst. Otherwise, the interlayer spacing of the swollen clay is sufficient to receive the Al and Al/Fe polyoxocations [1]. Because of that, bentonite has to be modified to get more stabile structure. One of the ways to modify bentonite is by pillarization process [2]. The application fields of these substrates are very broad because of the possibility to intercalate different types of pillars making them suitable for specific adsorption and catalytic processes.

The introduction of inorganic pillars in the natural bentonite improves its resistance and stability, increases its micro porosity and provides larger surface area and accessibility to its acid sites (Brönsted and Lewis sites) [3],[4]. Pillared bentonite can be utilized as catalyst because of the reason that pillared bentonite has active sites such as silica group (SiO_4) and alumina group (AlO_4). The presence of active metal which is penetrated into bentonite structure during pillarization process can enhance the catalytic performance of pillared bentonite. Other wise, the presence of pore which has definite size will give the ability to select both of reactants and products.

Various works have been reported on the catalyst synthesis. They show that catalytic activity depends on several factors such as chemical composition, preparation method, calcinations temperature and characteristic of catalyst surface.

In environmental catalysis there is a great interest in obtaining pillared clays containing iron (oxides or oxyhydroxides) species use in the catalytic wet peroxide oxidation of toxic organic compounds since many industrial processes yield a variety of organic contaminants which are poured into natural water sources with negative impact on ecosystem and humans (toxicity, carcinogenic and mutant properties) [5]. Among these pollutants contained in industrial residual water we can highlight phenol, substituted phenols (chlorophenol, nitrophenol), oxalic acid, acetic acid, pesticides and herbicides [5]-[8]. Phenol is particularly considered as one of the most toxic organic contaminants and it is commonly chosen as model molecule on catalytic oxidation of organic compounds studies in diluted aqueous medium [9].

This work shows the synthesis of solid catalyst, bentonite pillared with Al/Fe and intercalated with HDTMA, in which iron is introduced as active phase (activation sites for hydrogen peroxide as well as organic molecules) in order to develop the catalyst material to the complete oxidation of organic molecules.

Experimental

Catalyst preparation

The starting clay was a natural Pacitan bentonite, extracted from Pacitan region, East Java. The pillaring agent solution was prepared by mixing NaOH and Fe (and NaOH Al/Fe 1 mmole) which has molar ratio OH to Fe = 0.8 (and OH to Al/Fe = 0.8). The intercalating agent solution was prepared by mixing HDTMA 0.1% solution and bentonite suspension which has total volume of solution was 1 L. Pillarization process of the clay was carried out by mixing bentonite and pillaring agent solution with ratio [gram bentonite/volume of solution] = 1 gram/50 mL. The mixture was heated at 70-75 °C. After 5 hours, the mixture was cooled, and washed with aquadest. The obtained solid was dried in the oven at 100 °C. After pillarization process, bentonite was calcined at 500 °C for 4 hours with nitrogen and oxygen stream.

Characterization of the catalyst

The chemical analysis of the solids was carried out by Fourier Transform Infra Red Spectrophotometer (FTIR, Bruker Tensor 27, Germany) to characterize bentonite structure. The amount of Fe was analyzed using Genesys 6 UV-Vis Spectrophotometer.

Catalytic activity

The catalytic oxidation reaction of phenol in a diluted aqueous medium was carried out in a batch glass reactor equipped with thermometer, and reflux condenser. It is open to the atmosphere, and thoroughly stirring with magnetic stirrer. Firstly, the reactor was loaded with phenol solution and catalyst and heated up until 70°C. The reaction was conducted at 70°C for 4 hours. The hydrogen peroxide solution was added stepwise during a reaction time of 4 hours. The molar ratio of phenol to hydrogen peroxide = 1:1, weight ratio of catalyst to phenol = 1:10. The course of the phenol conversion and selectivity was followed by high performance liquid chromatography (HPLC) by means of Knauer.

Results and Discussion

Characterization

Fig.1. depicts FTIR spectra of the catalyst materials, i.e. fresh bentonite, Fe pillared bentonite with HDTMA 0,1 %, Fe pillared bentonite, Al-Fe pillared bentonite with HDTMA 0,1 % and AL-Fe pillared bentonite.




Fig. 1. FTIR Spectra of fresh bentonite, PILB-HDTMA 0,1 % Fe 1 mmole, PILB-Fe 1 mmole, PILB-HDTMA 0,1 % Al-Fe 1 mmole (1:1), PILB-Al-Fe 1 mmole (1:1), after calcination

The broad absorption peak at 3442 cm^{-1} indicates the presence of stretching asymmetric -OH group. It is supported by the absorption peak in the region 1638 cm^{-1} which is indicated stretching symmetric H-O-H of water. The absorption peak at 3626 cm^{-1} appears clearly for all of material because bentonite has naturally -OH group in the form of Al-OH , and Si-OH in the matrix polymer.

The absorption peaks in the region $1030 - 1040\text{ cm}^{-1}$ indicate asymmetric stretching O-Si-O and O-Al-O group in tetrahedral structure TO_4 ($\text{T} = \text{Si}$ or Al) of clay structure. The peak would be appeared wider after pillaring process with Fe because of the new formation of polymer system O-Al-O and O-Si-O in pillared bentonite. Figure 1 also shows the broad absorption peak in the range of $950 - 1200\text{ cm}^{-1}$ but not significant. The peak indicates asymmetric stretching of O-Fe-O in pillared bentonite structure. It reflects the presence of metal Fe which penetrates, forms pillars and supports layers in the bentonite structure as Fe oxide. However the spectra show that the number of Fe is not significant. It can be shown from table 1 that the amount of Fe which penetrates in each sample is not too different from it in the absorption peak in the range $950 - 1200\text{ cm}^{-1}$.

Table 1. The amount of Fe which penetrates in bentonite structure

Sample	The amount of Fe (mg/g bentonite)
PILB Fe 1 mmole	4,8088
PILB Al-Fe 1 mmole (1:1)	3,5147
PILB HDTMA 0,1% Fe 1 mmole	5,0147
PILB HDTMA 0,1% Al-Fe 1 mmole (1:1)	5,1029

The highest amount of Fe which penetrates into bentonite is PILB-HDTMA 0,1 % Al-Fe 1 mmole (1:1) and PILB HDTMA 0,1 % Fe 1 mmole. The result is validated by the IR result at figure 1 where the broadest absorption peak in the range 950 – 1200 cm⁻¹ is PILB HDTMA 0,1 % Al-Fe 1 mmole (1:1) and PILB HDTMA 0,1 % Fe 1 mmole. PILB HDTMA 0,1 % Al-Fe 1 mmole (1:1) and PILB HDTMA 0,1 % Fe 1 mmole have higher polymerization of O-Fe-O than PILB Al-Fe 1 mmole (1:1) and PILB Fe 1 mmole even though the difference is not significant.

The spectral region of 500 – 600 cm⁻¹ show the presence of double ring structure which indicates there are pores in the pillared bentonite. Unfortunately, the pore structure of pillared bentonite does not differ significantly from it of fresh bentonite. The fact shows that pillaring process has not given pores with sufficient size.

Catalytic activity test

Phenol hydroxylation is oxidation of phenol by using H₂O₂. Molar ratio of phenol to H₂O₂ is 1 : 1. Reaction was conducted in batch mode in the presence of pillared bentonite as solid catalyst. The weight ratio of catalyst to phenol was 1:10, and concentration of phenol is 1 M. The reaction temperature was 70°C, and reaction time was 4 hours. The observed variables were catalyst activity and selectivity. The catalyst activity was observed as phenol conversion, and the selectivity was observed as selectivity of wanted product (hydroquinone).

Table 2 Hydroxylation of phenol to hydroquinone and cathecol

Catalyst	Phenol Conversion (%)	Selectivity (%)		
		hydroquinone	cathecol	benzoquinone
Without catalyst	1.2	49.10	50.30	0.6
Fresh bentonite	4.4	47.00	52.40	0.6
PILB Fe 1 mmole	78.0	61.80	38.10	0.1
PILB Al-Fe 1 mmole (1:1)	72.7	59.77	40.20	0.03

PILB HDTMA 0,1% Fe 1 mmole	65.6	58.20	41.71	0.09
PILB HDTMA 0,1% Al-Fe 1 mmole (1:1)	76.7	63.37	36.60	0.03

Reaction condition: Catalyst/Phenol/H₂O₂ = 1/10/10

Percentage conversion and reaction selectivity were determined by HPLC analysis

In general, the presence of catalyst can increase reactant conversion and product selectivity. Table 2 shows phenol conversion and catalytic product selectivity (hydroquinone and cathecol) increase in the presence of catalyst. It is caused chemical adsorption in the catalyst pore surface will release energy so that it will decrease activation energy. Beside of the reason, Fe in the catalyst structure can oxidize phenol together with H₂O₂.

PILB Fe 1 mmole catalyst gives phenol conversion as high as 78% and the amount of Fe penetrates into bentonite is 4.8 mg/g solid. Whereas PILB HDTMA 0.1 % Fe 1 mmole catalyst reach phenol conversion 65.6 % and the amount of Fe introduces into solid is 5 mg/g solid. The amount of Fe which is introduced into PILB HDTMA 0.1 % Fe 1 mmole is slightly higher than that of PILB Fe 1 mmole catalyst but the phenol conversion of PILB HDTMA 0.1 % Fe 1 mmole is lower than that of PILB Fe 1 mmole. It is caused that phenol has been converted into catalytic product in the system with PILB HDTMA 0.1 % Fe 1 mmole. The selectivity of hydroquinone and cathecol of system which uses PILB HDTMA 0.1 % Fe 1 mmole in series are 58.2% and 41.7%, whereas those of PILB Fe 1mmole catalyst are 61.8% and 38.1%. This result describes that HDTMA molecule on PILB HDTMA 0.1 % Fe 1 mmole has not been very successfully disappeared during calcinations and oxidation process, since the selectivity of wanted product on PILB HDTMA 0.1 % Fe 1 mmole is lower than that of on PILB Fe 1mmole. This is supported by FTIR spectra which is the pore of PILB HDTMA 0.1 % Fe 1 mmole has not significantly created.

All modified bentonite catalyst will give better result (higher phenol conversion and CAT selectivity) compare to fresh bentonite. It shows that modification process (intercalation and pillarisation) give slightly effect to increase catalytic activity of fresh bentonite because of increasing bentonite pore size.

Based on this finding, it is proposed that Fe has participated in this reaction. HDTMA is expected to increase pore size and be able to support two bentonite layers before penetrating pillaring agent, but the experiment did not get optimal condition of material. It is proved by reaction products (HQ and CAT) which is produced by using catalyst with HDTMA is the same as that which is produced by using catalyst without HDTMA. It is

caused HDTMA can not be disappeared and give crowded pores in the bentonite structure. Because of that the reaction can not run effectively, because HDTMA has covered active sites of catalyst. This condition causes small amount of reactant can be contacted with active site of catalyst so that the reaction tends to follow non catalytic mechanism. In this study the presence of Al does not give significant increase in the activity of catalyst.

Conclusion

In this research, pilllarization process of bentonite using Fe has successfully given significant increase in the pore size of natural bentonite so that it increase the catalytic activity and product selectivity. The presence of Al and HDTMA in intercalation process does not give very significant effect to increase pore size and catalytic activity to the pillared-Fe bentonite.

Acknowledgment

The authors offer their gratitude to Expertise Area of Development Program (EADP) of University of Surabaya for the financial support of the project.

References

1. L. Storaro, M. Lenarda, R. Ganzerla, and A. Rinaldi, *Microp Mesop Mater* 6 (1996), 55-63
2. Deng, Y., Dixon, J.B. And G. White, G. N., *Clays Clay Miner.* 51 (2004), 150–161
3. L. Bergaoui, J.F. Lambert, R. Franck, H. suquet, J.L., Robert, *J.Chem. Soc.* 91 (1995), 2229
4. J.M. Thomas, W.J., Thomas, “Principles and Practice of Heterogeneous Catalysis, VCH Verlag, Weinheim, 1997
5. Davis, R.D., Gilman, J.W., Sutto, T.E., Callahan, J.H., Trulove, P.C. And De Long, H.C., *Clays Clay Miner* 52 (2004), 171–179.
6. He. H., Frost, R.L., Deng, F., Zhu, J., Wen, X. And Yuan, P., *Clays Clay Miner* 52 (2004), 350–356
7. Slade, P.G., And Gates, W.P., *Clays Clay Miner* 52 (2004), 204-210.
8. Weiss, Z., Valášková, M., Křístková, M., Čapková, P., Pospíšil, M., *Clays Clay Miner* 51 (2004), 555-565.
9. R. K. Widi, Arief B., Lieke R., *Inter J. Appl. Chem.* 6 (2009) 1, 11-18